COLORADO ASTHMA CARE PLAN AND MEDICATION ORDER FOR SCHOOL AND CHILD CARE SETTINGS*

PARENT/GUARDIAN COMPLETE, SIGN AND DATE:			
Child Name:			Birthdate:
School:			Grade:
Parent/Guardian Name: Phone:			
I approve this care plan and give permission for school personnel to share this information, follow this plan, administer medication and care for my child/youth, and if necessary, contact our health care provider. I assume responsibility for providing the school/ program prescribed, non-expired medication and supplies (such as a spacer), and to comply with board policies, if applicable. I am aware <i>911 may be called if a quick relief inhaler is not at school</i> and my child/youth is experiencing symptoms.			
Parent/Guardian Signature			Date
HEALTH CARE PROVIDER COMPLETE ALL ITEMS, SIGN AND DATE:			
QUICK RELIEF MEDICATION: Albuterol Other:			
Common side effects: ↑ heart rate, tremor □ Use spacer with inhaler (MDI) Controller medication used at home:			
TRIGGERS: ☐ Weather ☐ Illness ☐ Exercise ☐ Smoke ☐ Dust ☐ Pollen ☐ Poor Air Quality ☐ Other:			
☐ Life threatening allergy specify:			
QUICK RELIEF INHALER ADMINISTRATION: With assistance or self-carry.			
Student needs supervision or assistance to use inhaler. Student will not self-carry inhaler.			
Student understands proper use of asthma medications, and in my opinion, can self-carry and use his/her inhaler at school independently with approval from school nurse and completion of contract.			
IF YOU SEE THIS: DO THIS:			DO THIS:
GREEN ZONE: No Symptoms Pretreat	No current symptomsStrenuous activity	☐ Not required <u>OR</u> ☐ Student	US ACTIVITY , please choose ONE : /Parent request <u>OR</u> □ Routinely
	planned		ninutes before activity: 2 puffs 4 puffs
		Repeat in 4 hours, if needed for	additional physical activity. cing symptoms, follow YELLOW or RED ZONE.
	Trouble breathing	1. Give QUICK RELIEF MED: 2 puffs 4 puffs	
NE:	• Wheezing	2. Stay with child/youth and maintain sitting position.	
YELLOW ZONE: Mild symptoms	 Frequent cough 	3. REPEAT QUICK RELIEF MED if not improving in 15 minutes: \square 2 puffs \square 4 puffs	
	Chest tightness	If symptoms do not improve or worsen, follow RED ZONE.	
	 Not able to do activities 	4. Child/youth may go back to normal activities, once symptoms are relieved.5. Notify parents/guardians and school nurse.	
5. Notify parents/guardians and school harse.			
ns Ins	Struggles to breathe	 Give QUICK RELIEF MED: 2 puffs 4 puffs Refer to the anaphylaxis care plan if the student has a life threatening allergy. If there is no anaphylaxis care plan follow emergency guidelines for anaphylaxis. 	
.: 스 bi	 Trouble talking (only 		
RED ZONE: EMERGENCY Severe Symptoms	speaks 3-5 words)	2. Call 911 and inform EMS the reason for the call.	
	Skin of chest and/or neck Skin of chest and/or neck	 REPEAT QUICK RELIEF MED if not improving: □ 2 puffs □ 4 puffs Can repeat every 5-15 minutes until EMS arrives. Stay with child/youth. Remain calm, encouraging slower, deeper breaths. 	
	pull in with breathingLips/fingernails gray/blue		
Š	- Lips/ inigeritans gray/ blac	5. Notify parents/guardians and school nurse.	
Health Care Provider Signature Print Provider Name Date			
Good for 12 months unless specified otherwise in district policy.			
Fax Pho		one	Email
School Nurse/CCHC Signature Date			
☐ Self-carry contract on file. ☐ Anaphylaxis plan on file for life threatening allergy to:			

^{*}Including reactive airways, exercise-induced bronchospasm, twitchy airways.

